

## LIST OF FIGURES

Figure	Page
2-1 Basins in the MBSS study area and the years scheduled for sampling in the 1995-1997 survey . . . . .	2-2
2-2 MBSS stratified random sampling design . . . . .	2-4
2-3 Randomly selected sites sampled in the 1995-1997 MBSS . . . . .	2-6
3-1 Precipitation isobar, annual runoff per square mile of watershed area, and mean annual temperature isobar maps for Maryland . . . . .	3-2
3-2 Surface elevation map of Maryland (Adapted from MGS 1996) . . . . .	3-4
3-3 Lithogeochemistry map of Maryland . . . . .	3-6
3-4 Soil erodibility map of Maryland . . . . .	3-10
3-5 Land use map of Maryland . . . . .	3-11
3-6 Road network map of Maryland . . . . .	3-13
3-7 Population density map of Maryland based on 1990 census . . . . .	3-14
3-8 Airshed and watershed map for the Chesapeake Bay . . . . .	3-15
3-9 Map of dams and other barriers to fish migration in Maryland . . . . .	3-17
4-1 Per-segment fish species richness (mean number of species per 75-m segment), statewide and for basins sampled in the 1995-1997 MBSS . . . . .	4-6
4-2 Per-segment fish species richness (mean number of species per 75-m segment), by stream order, for the 1995-1997 MBSS . . . . .	4-7
4-3 Fish density (number of individuals per stream mile), statewide and for basins sampled in the 1995-1997 MBSS . . . . .	4-7
4-4 Fish density (number of individuals per stream mile) by stream order, for the 1995-1997 MBSS. Error bars signify $\pm 1$ standard error (lack of error bars indicate that variance is statistically undefined) . . . . .	4-9
4-5 Estimated percentage of stream miles with no fish, statewide and for basins sampled in the 1995-1997 MBSS . . . . .	4-11
4-6 Fish biomass (kg per stream mile), statewide and for basins sampled in the 1995-1997 MBSS . . . . .	4-12
4-7 Density (number of individuals per stream mile) of brook trout ( <i>Salvelinus fontinalis</i> ), statewide and for the basins sampled in the 1995-1997 MBSS . . . . .	4-14
4-8 Density (number of individuals per stream mile) of brook trout ( <i>Salvelinus fontinalis</i> ), by stream order for the basins sampled in the 1995-1997 MBSS . . . . .	4-15
4-9 Total gamefish density (number of individuals per stream mile), statewide and for basins sampled in the 1995-1997 MBSS . . . . .	4-16
4-10 Total gamefish density (number of individuals per stream mile), by stream order for the 1995-1997 MBSS . . . . .	4-17
4-11 Gamefish biomass (kg per stream mile), statewide and for basins sampled in the 1995-1997 MBSS . . . . .	4-18
4-12 Estimates of the total abundance of harvestable size gamefish (number of individuals), statewide and for basins sampled in the 1995-1997 MBSS . . . . .	4-19
4-13 Percentage of fish with each type of pathological anomaly (skin, skeletal, or ocular), statewide for the 1995-1997 MBSS . . . . .	4-28
4-14 Occurrence of pathological anomalies in gamefish (percent of individual gamefish with pathological anomalies), statewide and for basins sampled in the 1995-1997 MBSS . . . . .	4-30
4-15 Occurrence of pathological anomalies in nongame fish (percent of nongame fish with pathological anomalies), statewide and for basins sampled in the 1995-1997 MBSS . . . . .	4-32
4-16 Percentage of stream miles containing fish with each type of pathological anomaly (skin, skeletal or ocular), statewide for the 1995-1997 MBSS . . . . .	4-33
4-17 Percentage of stream miles with gamefish species having pathological anomalies, statewide and for basins sampled in the 1995-1997 MBSS . . . . .	4-35

## LIST OF FIGURES (Cont'd)

Figure	Page
4-18 Percentage of stream miles with nongame fish species having pathological anomalies, statewide and for basins sampled in the 1995-1997 MBSS .....	4-37
4-19 Number of benthic taxa, statewide and by basin for the 1995-1997 MBSS .....	4-39
4-20 Mean benthic taxa richness (mean number of benthic taxa per site), statewide and for basins sampled in the 1995-1997 MBSS .....	4-41
4-21 Mean benthic taxa richness (mean number of benthic taxa per site), statewide, by watershed size .....	4-42
4-22 Percentage of stream miles with salamanders present, by stream order for the 1995-1997 MBSS .....	4-44
4-23 Percentage of stream miles with salamanders, statewide and for basins sampled in the 1995-1997 MBSS .....	4-45
4-24 Percentage of stream miles with frogs, statewide and for basins sampled in the 1995-1997 MBSS .....	4-46
4-25 Number of mussel species, statewide and for basins sampled in the 1995-1997 MBSS .....	4-48
4-26 Distribution of native and non-native mussels species recorded in the 1995-1997 MBSS .....	4-49
4-27 Mean number of aquatic plant species per site based on the percent shading received at each site for the 1995-1997 MBSS .....	4-51
4-28 Distribution of submerged aquatic vegetation recorded the 1995-1997 MBSS .....	4-52
4-29 Mean number of aquatic plant species per site, by stream order, for Coastal Plain and non-Coastal Plain sites in the 1995-1997 MBSS .....	4-53
5-1 The three geographic regions used for the derivation of the fish Index of Biotic Integrity: Coastal, Piedmont, and Highland .....	5-2
5-2 Derivation and interpretation of scores for the MBSS fish and benthic Indices of Biotic Integrity .....	5-8
5-3 Geographic distribution of fish Index of Biotic Integrity scores for basins sampled in the 1995-97 MBSS .....	5-11
5-4 Fish Index of Biotic Integrity (IBI) scores for basins sampled in the 1995-97 MBSS .....	5-12
5-5 Fish Index of Biotic Integrity (IBI) scores by stream order, for basins sampled in the 1995-97 MBSS .....	5-13
5-6 Geographic distribution of fish Index of Biotic Integrity (IBI) scores throughout the study area, including the statewide distribution of the percentage of stream miles with fish in each category .....	5-15
5-7 Geographic distribution of benthic Index of Biotic Integrity scores for basins sampled in the 1995-97 MBSS .....	5-16
5-8 Benthic Index of Biotic Integrity (IBI) scores for basins sampled in the 1995-1997 MBSS .....	5-17
5-9 Benthic Index of Biotic Integrity (IBI) scores by stream order, for basins sampled in the 1995-97 MBSS .....	5-18
5-10 Geographic distribution of benthic Index of Biotic Integrity (IBI) scores throughout the study area, including the statewide distribution of the percentage of stream miles with benthic IBI scores in each category .....	5-20
5-11 Hilsenhoff Biotic Index scores for basins sampled in the 1995-97 MBSS .....	5-22
5-12 Hilsenhoff Biotic Index scores by stream order, for basins sampled in the 1995-97 MBSS .....	5-23
5-13 Relationship between fish IBI and benthic IBI for the Patapsco basin .....	5-24
5-14 Relationship between benthic IBI and Hilsenhoff Biotic Index for the Patuxent basin .....	5-24
6-1 Percentage of stream miles with low pH by basin (spring pH), for basins sampled in the 1995-1997 MBSS .....	6-3
6-2 Percentage of stream miles with low pH by basin (summer pH), for basins sampled in the 1995-1997 MBSS .....	6-4
6-3 Percentage of stream miles with low ANC by basin, for basins sampled in the 1995-1997 MBSS .....	6-6
6-4 Procedure for the determination of acid sources for sites sampled in the 1995-1997 MBSS .....	6-7
6-5 Relationship between nitrate-nitrogen (NO <sub>3</sub> -N) and the percentage of agricultural land use for the basins sampled in the 1995-1997 MBSS .....	6-9
6-6 Percentage of stream miles with ANC < 200 µeq/l, by acid source for the basins sampled in the 1995-1997 MBSS .....	6-10
6-7 Acid sources at 1995-1997 MBSS sites .....	6-11
6-8 Percentage of stream miles with ANC < 200 µeq/l by acid source, by stream order for the 1995-1997 MBSS ..	6-13
6-9 Statewide percentage of stream miles with ANC < 200 µeq/l and with ANC < 0 µeq/l by acid source .....	6-14

## LIST OF FIGURES (Cont'd)

Figure	Page
6-10 Fish IBI by summer pH class (< 5, 5-6, > 6) for the 1995-1997 MBSS .....	6-17
6-11 Fish and amphibian and reptile species richness by ANC class (< 0, 0-50, 50-200, > 200 $\mu\text{eq/l}$ ) for the 1995-1997 MBSS .....	6-18
6-12 Statewide biomass estimates (kg/stream mile) for nongame fish and gamefish by ANC class(< 0, 0-50, 50-200, > 200 $\mu\text{eq/l}$ ), 1995-1997 MBSS .....	6-18
6-13 Benthic IBI by spring pH class (< 5, 5-6, > 6) for the 1995-1997 MBSS .....	6-19
6-14 Fish and benthic IBI by ANC class (< 0, 0-50, 50-200, > 200 $\mu\text{eq/l}$ ) for the 1995-1997 MBSS .....	6-19
7-1 Percentage of stream miles by riparian buffer type and width for the 1995-1997 MBSS .....	7-3
7-2 Riparian buffer width at sites sampled in the 1995-1997 MBSS .....	7-4
7-3 Percentage of stream miles with riparian buffer width less than 5 meters, statewide and for the basins sampled in the 1995-1997 MBSS .....	7-5
7-4 Percentage of stream miles with beaver ponds, statewide and for the basins sampled in the 1995-1997 MBSS ..	7-6
7-5 Percentage of stream miles with evidence of channelization, statewide and for the basins sampled in the 1995-1997 MBSS .....	7-8
7-6 Percentage of stream miles in "Highest" and "High" categories of the bank erodibility index, statewide and for the basins sampled in the 1995-1997 MBSS .....	7-9
7-7 Percentage of stream miles with poor and marginal instream habitat structure, statewide and for the basins sampled in the 1995-1997 MBSS .....	7-10
7-8 Percentage of stream miles with poor and marginal epifaunal substrate, statewide and for the basins sampled in the 1995-1997 MBSS .....	7-11
7-9 Percentage of stream miles with poor and marginal velocity/depth diversity, statewide and for the basins sampled in the 1995-1997 MBSS .....	7-12
7-10 Percentage of stream miles with poor and marginal pool/glide/eddy quality, statewide and for the basins sampled in the 1995-1997 MBSS .....	7-13
7-11 Percentage of stream miles with poor and marginal riffle/run quality, statewide and for the basins sampled in the 1995-1997 MBSS .....	7-14
7-12 Percentage of stream miles with poor and marginal aesthetic quality, statewide and for the basins sampled in the 1995-1997 MBSS .....	7-17
7-13 Percentage of stream miles rated as easy and moderately easy access, statewide and for the basins sampled in the 1995-1997 MBSS .....	7-18
7-14 Relationship between aesthetic quality and remoteness, statewide, for the 1995-1997 MBSS .....	7-19
7-15 Mean stream width, statewide and for the basins sampled in the 1995-1997 MBSS .....	7-20
7-16 Mean thalweg depth, statewide and for the basins sampled in the 1995-1997 MBSS .....	7-21
7-17 Mean discharge, statewide and for the basins sampled in the 1995-1997 MBSS .....	7-22
7-18 Mean number of pieces of wood found in the stream, statewide and for the basins sampled in the 1995-1997 MBSS .....	7-23
7-19 Geographic distribution of Physical Habitat Index (PHI) ratings for sites sampled in the 1995-1997 MBSS ...	7-26
7-20 Physical Habitat Index (PHI) ratings for the basins sampled in the 1995-1997 MBSS, as the percentage of stream miles in each category .....	7-27
7-21 Distribution of Physical Habitat Index (PHI) ratings for the basins sampled in the 1995-1997 MBSS as the percentage of stream miles in each category .....	7-29
7-22 Relationship between the fish IBI and the Physical Habitat Index (PHI), statewide for the 1995-1997 MBSS ..	7-32
7-23 Relationship between the benthic IBI and the Physical Habitat Index (PHI), statewide for the 1995-1997 MBSS .....	7-32
7-24 Mean number of amphibian and reptile species in three categories of the Physical Habitat Index (PHI) for the 1995-1997 MBSS .....	7-33

## LIST OF FIGURES (Cont'd)

Figure	Page
7-25 Relationship between the fish IBI and instream habitat structure, statewide for the 1995-1997 MBSS . . . . .	7-33
7-26 Relationship between the fish IBI and aesthetic quality, statewide for the 1995-1997 MBSS . . . . .	7-34
7-27 Relationship between the fish IBI and maximum depth, statewide for the 1995-1997 MBSS . . . . .	7-34
7-28 Relationship between the benthic IBI and riffle/run quality, statewide for the 1995-1997 MBSS . . . . .	7-35
7-29 Relationship between the benthic IBI and aesthetic quality, statewide for the 1995-1997 MBSS . . . . .	7-35
7-30 Relationship between the fish IBI and local riparian buffer width, statewide for the 1995-1997 MBSS . . . . .	7-36
7-31 Relationship between the benthic IBI and epifaunal substrate, statewide for the 1995-1997 MBSS . . . . .	7-36
7-32 Mean number of amphibian and reptile species per site for each category of riparian buffer width, statewide, for the 1995-1997 MBSS . . . . .	7-37
8-1 Nitrate nitrogen (NO <sub>3</sub> -N) concentration (mg/l) statewide and for basins sampled in the 1995-1997 MBSS . . . . .	8-3
8-2 Mean nitrate nitrogen (NO <sub>3</sub> -N) concentration (mg/l) statewide and for basins sampled in the 1995-1997 MBSS . . . . .	8-4
8-3 Mean number of benthic Ephemeroptera, Plecoptera, and Trichoptera (EPT) taxa declined with higher nitrate nitrogen concentration at 1995-1997 MBSS sites . . . . .	8-5
8-4 Relationship between nitrate nitrogen (NO <sub>3</sub> -N) concentration (mg/l) and the Hilsenhoff Biotic Index for the 1995-1997 MBSS . . . . .	8-5
8-5 Dissolved oxygen (DO) concentration (ppm) statewide and for basins sampled in the 1995-1997 MBSS . . . . .	8-6
8-6 Mean nitrate nitrogen (NO <sub>3</sub> -N) concentration (mg/l) for CORE/Trend stations sampled in April and May of 1995, 1996, and 1997 . . . . .	8-7
8-7 Mean nitrate nitrogen (NO <sub>3</sub> -N) concentration (mg/l) for CORE/Trend stations sampled in April and May of 1995-1997 and for MBSS sites sampled during the spring index period of 1995-1997 . . . . .	8-8
9-1 Major land use types within individual catchments upstream of the 1995-1997 MBSS sampling sites . . . . .	9-8
9-2 Percentage of three land use types (forest, urban, and agriculture) for two streams in the Patuxent basin - Dorsey Run and Midway Run . . . . .	9-9
9-3 Water temperature (°C) during July 1997 for two streams in the Patuxent basin - Dorsey Run and Midway Run . . . . .	9-10
9-4 Mean nitrate-nitrogen concentration (mg/l), statewide and for basins sampled in the 1995-1997 MBSS, among sites with catchment land use less than and greater than 50% agriculture . . . . .	9-11
9-5 Relationship between the fish IBI and urban land use for the basins sampled in the 1995-1997 MBSS . . . . .	9-12
9-6 Relationship between the fish IBI and urban land use for the Potomac Washington Metro basin . . . . .	9-13
9-7 Relationship between the fish IBI and urban land use for the Patapsco basin . . . . .	9-14
9-8 Relationship between the fish IBI and low-intensity development for the basins sampled in the 1995-1997 MBSS . . . . .	9-15
9-9 Relationship between the fish IBI and agricultural land use for the basins sampled in the 1995-1997 MBSS . . . . .	9-16
9-10 Relationship between the fish IBI and agricultural land use for the Gunpowder basin . . . . .	9-17
9-11 Relationship between the fish IBI and forested land cover for the basins sampled in the 1995-1997 MBSS . . . . .	9-18
9-12 Relationship between the benthic IBI and urban land use for the basins sampled in the 1995-1997 MBSS . . . . .	9-19
9-13 Relationship between the benthic IBI and urban land use for the Potomac Washington Metro basin . . . . .	9-20
9-14 Relationship between the benthic IBI and urban land use for the Patuxent basin . . . . .	9-21
9-15 Relationship between the benthic IBI and low-intensity development for the basins sampled in the 1995-1997 MBSS . . . . .	9-22
9-16 Relationship between the benthic IBI and agricultural land use for the basins sampled in the 1995-1997 MBSS . . . . .	9-23

## LIST OF FIGURES (Cont'd)

Figure	Page
9-17 Relationship between the benthic IBI and forested land cover for the basins sampled in the 1995-1997 MBSS .....	9-24
9-18 Relationship between the Hilsenhoff Biotic Index and urban land use for the basins sampled in the 1995-1997 MBSS .....	9-25
9-19 Relationship between the Hilsenhoff Biotic Index and urban land use for the Potomac Washington Metro basin .....	9-26
9-20 Relationship between the Hilsenhoff Biotic Index and low-density development for the basins sampled in the 1995-1997 MBSS .....	9-27
9-21 Relationship between the Hilsenhoff Biotic Index and agricultural land use for the basins sampled in the 1995-1997 MBSS .....	9-28
9-22 Relationship between the Hilsenhoff Biotic Index and forested land cover for the basins sampled in the 1995-1997 MBSS .....	9-29
9-23 Relationship between the number of aquatic salamanders per site and urban land use for the basins sampled in the 1995-1997 MBSS .....	9-30
10-1 Statewide percent deviation from normal precipitation amount for the MBSS sample years 1995-1997 .....	10-3
10-2 Mean discharge (cfs) for the three basins that were sampled in multiple years of the 1995-1997 MBSS .....	10-4
10-3 Mean fish Index of Biotic Integrity (IBI) scores for the three basins that were sampled in multiple years of the 1995-1997 MBSS .....	10-5
10-4 Mean benthic Index of Biotic Integrity (IBI) scores for the three basins that were sampled in multiple years of the 1995-1997 MBSS .....	10-6
10-5 Mean Physical Habitat Indicator scores for the three basins that were sampled in multiple years of the 1995-1997 MBSS .....	10-7
10-6 Mean nitrate nitrogen (NO <sub>3</sub> -N) concentrations (mg/l) for the three basins that were sampled in multiple years of the 1995-1997 MBSS .....	10-8
11-1 Comparative ranking of stressors affecting streams in the 1995-1997 MBSS .....	11-2
11-2 Extent of stressors affecting streams for basins sampled in the 1995-1997 MBSS .....	11-3
12-1 Native fish species richness by site for the 1995 - 1997 MBSS .....	12-3
12-2 Fish species richness by basin and geographic region for the 1995-1997 MBSS .....	12-4
12-3 Benthic species richness by basin and geographic region for the 1995-1997 MBSS .....	12-5
12-4 Terrestrial and aquatic amphibian and reptile species richness by basin for the 1995-1997 MBSS .....	12-7
12-5 Total amphibian and reptile species richness by basin and geographic region for the 1995-1997 MBSS .....	12-8
12-6 Salamander species richness by basin and geographic region for the 1995-1997 MBSS .....	12-9
12-7 Frog and toad species richness by basin and geographic region for the 1995-1997 MBSS .....	12-10
12-8 Turtle species richness by basin and geographic region for the 1995-1997 MBSS .....	12-11
12-9 Snake and lizard species richness by basin and geographic region for the 1995-1997 MBSS .....	12-12
12-10 Mussel species richness by basin and geographic region for the 1995-1997 MBSS .....	12-14
12-11 Aquatic plant species richness by basin and geographic region for the 1995-1997 MBSS .....	12-15
12-12 Distribution of state-listed rare and endangered fish found in the 1995 - 1997 MBSS, by watershed .....	12-16
12-13 Overlay of watershed fish species richness and sites with rare fish species .....	12-18
12-14 Presence of Asiatic clam ( <i>Corbicula fluminea</i> ) at 1995-1997 MBSS sampling sites .....	12-22
12-15 Density of native and non-native fish species for basins sampled in the 1995-1997 MBSS .....	12-23
12-16 Density of non-native fish species for basins sampled in the 1996-1997 MBSS .....	12-25
12-17 Percentage of stream miles with non-native fish species, by stream order, 1995-1997 MBSS .....	12-26
12-18 Sites that rated good (IBI scores ≥ 4.0) for both the fish and benthic Index of Biotic Integrity .....	12-28

## LIST OF FIGURES (Cont'd)

Figure	Page
13-1 Maryland's 10 Tributary; Strategies basins in the 1995-1997 MBSS .....	13-11
13-2 Maryland's 138 8-digit watersheds (in red) within the major river basins used in MBSS reporting (blue) ...	13-11
13-3 Percentage of stream miles with riparian buffer width 19-50 m, statewide and for the basins sampled in the 1995-1997 MBSS .....	13-13
13-4 Percentage of stream miles that are located on public, private, or mixed ownership land, statewide and for the basins sampled in the 1995-1997 MBSS .....	13-14
13-5 Geographic distribution of 1995-1997 MBSS sites that are located on private, public, or mixed ownership land .....	13-15
13-6 Geographic distribution of protected lands .....	13-16